

RESEARCH ARTICLE

Open Access

Plants accumulating heavy metals in the Danube River wetlands

Marius L Matache^{1*}, Constantin Marin², Laurentiu Rozylowicz¹ and Alin Tudorache²

Abstract

Background: We present herein our results regarding the accumulation of four heavy metals (copper, cadmium, lead, and zinc) in four aquatic species plants (*Ceratophyllum demersum*, *Potamogeton pectinatus*, *Potamogeton lucens*, *Potamogeton perfoliatus*) collected from the Danube River, South-Western part of Romania and their possible use as indicators of aquatic ecosystems pollution with heavy metals.

Methods: Elements concentration from the vegetal material was determined through Inductively Coupled Plasma – Mass Spectrometry.

Results: The species were chosen based on their previous use as bioindicators in aquatic ecosystems and due to the fact they are one of the most frequent aquatic plant species of the Danube River ecosystems within the Iron Gates Natural Park. Highest amounts are recorded for *Ceratophyllum demersum* (3.52 µg/g for Cd; 22.71 µg/g for Cu; 20.06 µg/g for Pb; 104.23 µg/g for Zn). Among the *Potamogeton* species, the highest amounts of heavy metals are recorded in *Potamogeton perfoliatus* (1.88 µg/g for Cd; 13.14 µg/g for Cu; 13.32 µg/g for Pb; 57.96 µg/g for Zn). The sequence for the bioconcentration factors (BCFs) calculated in order to describe the accumulation of the four metals is Cd >> Zn > Pb > Cu. Increase of the zinc concentration determines an increase of the cadmium concentration (Spearman rho=0.40, p=0.02).

Conclusions: Despite the low ambiental levels of heavy metals, the four aquatic plants have the ability to accumulate significant amounts, which make them useful as biological indicators. BCF value for *Ceratophyllum demersum* indicated this species as a cadmium hyperaccumulator.

Background

Macrophytes have an important ecological role in aquatic ecosystems due to the multiple functions they play in trace elements cycles such as metals removal from water column, suspended matter, riverbed sediments and metals transfer to the food chains [1]. Moreover, the chemical reactions that the plants roots induce in the sediments they grow in increase the sediments capacity to uptake metals [2].

The phytoplankton shows a high accumulation rate of heavy metals entering the aquatic ecosystems [3,4]. Trace elements concentration in plants sometimes reach levels of concentration of six orders of magnitude higher than the environmental one [5]. As a consequence, a number of certain species were found to be useful indicators of the

water quality [6] or as solutions for wastewater treatment through phytoremediation processes [7,8], because native plants have the advantage of adapting to the stress conditions of the target pollutants [9].

Metal uptake by vegetation can be element-, plant species- [10,11] and plant tissue- specific [12,13]. For example, *Ceratophyllum demersum* was identified as a hyperaccumulator species for cadmium [14], *Salvinia minima* has the capacity to remove lead from aqueous solutions [15], whilst *Cladophora sp.* is able to hyperaccumulate arsenic [16]. The capacity of plants to hyperaccumulate trace elements is influenced by the presence of humic substances or other chelating substances [17,18], temperature and salinity [19] or amount of metal released through leaves [20].

The Danube River is a transboundary water body suffering a high anthropogenic pressure along its course [21] and a peak of sediments contamination with heavy metals in the Iron Gates area [22].

* Correspondence: media@portiledefier.ro

¹University of Bucharest, Centre for Environmental Research and Impact Studies, Bucharest, Romania

Full list of author information is available at the end of the article

Little information is available on the amount of heavy metals in aquatic plant species from the Danube River ecosystems in Romania and is limited to the Danube Delta [23,24]. A broad study that includes our target area of interest was performed on the Serbian side of the Danube River with *Ceratophyllum demersum* and *Potamogeton* genus samples collected from 36 sites on a 573 km length sector [25].

Detailed studies regarding the heavy metals levels of concentration in sediments have been performed on the analysed sector, since significant pollution sources are located in the area for more than four decades [26]. They represented the starting point in our survey for targeting certain elements. Our research aims are: 1) to investigate the amount of heavy metals in water and four plant species (*Ceratophyllum demersum*, *Potamogeton pectinatus*, *Potamogeton perfoliatus*, *Potamogeton lucens*) collected from the Danube River wetlands within the Iron Gates Natural Park in South-Western Romania and 2) to discuss the opportunity of using these aforementioned plant species as indicators of the aquatic ecosystems pollution level with heavy metals.

Methods

We collected water samples from 3 locations (Bazias, Divici, Coronini) in the analysed area, performing four sampling campaigns, one corresponding to each season. Samples were collected in high density polypropylene bottles and nitric acid was immediately added after collection to avoid analytes loss.

A number of 35 plant samples were collected between December 2010 and July 2012, during different vegetation seasons, in order to obtain accurate data set. The plant species under consideration were *Ceratophyllum demersum* (N=12 samples), *Potamogeton perfoliatus* (N=14), *Potamogeton lucens* (N=2) and *Potamogeton pectinatus* (N=7). For the *Potamogeton* genus, the collected sample was composed mainly from leaves and stems, but not roots. The epiphytic plants were removed before storing the samples in plastic bags and frozen at -20°C prior to chemical analysis. For each location, a sample was composed of three plants of the same species collected and pooled into a uniform sample. Samples were superficially washed with distilled water in order to remove particulate matters and then rinsed with ultrapure water (TKA Ultra Pure System GenPure, resistivity $18.2\text{ M}\Omega\times\text{cm}$).

The samples were dried at 105°C for 72 hours and then finely grinded prior to digestion. Sample digestion (approximately 2 grams) was performed in an open system with 8 mL of ultrapure concentrated HNO_3 in a Teflon jar and on an electric plate until the complete dissolution of the organic matrix. The remaining acid extract was diluted with 5 mL of ultrapure water and the

resulting solution was filtered on a filter paper with medium porosity (Whatman no 54) and collected in a 25 mL volumetric flask that was brought to constant volume with ultrapure water.

We have determined the elements concentration from the vegetal material using a NexION 300S (PerkinElmer, Shelton, CT, USA) ICP-MS instrument (Serial No. 81SN2032001), equipped with a S10 Autosampler. The operating parameters of the instrument are summarized in Table 1. All the solutions were prepared with ultrapure water (LaboStar™ TWF UV7 Ultrapure Water System, electric resistance $18.2\text{ M}\Omega\times\text{cm}$).

The calibration curves (Figure 1) were determined starting from a Multi Element Calibration Standard 3 solution provided by PerkinElmer (Part No. N9300233, Lot No. CL7-173YPY1). The solution contains 30 elements, each of them in a concentration of 10 mg/L. The working solutions for internal standard were prepared starting from the Internal Standard Mix, PerkinElmer Part No. N9303833, Lot No. CL43-46AS, (10 mg/L, Bi, Ge, In, ^6Li , Sc, Tb, Y) solution. Quality Control (QC), checking and Continuing Calibration Verification (CCV) solutions preparation used CertiPUR® Reference Material, ICP Multielement Standard Solution XXI (Merck, Cat. No. 109498, Lot No. HC 142053). HNO_3 60% Ultrapur® (Merck, Cat. No. 101518) was used for the vegetal material digestion, as well as the blanks preparation and washing solutions.

The solutions resulted from samples digestion were diluted 1:5, and the internal standard was added to the samples in order for each used element to reach a concentration of 20 $\mu\text{g/L}$. These were ^{74}Ge for ^{63}Cu and ^{66}Zn ; ^{115}In for ^{111}Cd ; ^{209}Bi for ^{208}Pb . The instrument was set to work for all elements in kinetic energy discrimination (KED) mode, with He gas flow rate of 3.6 and 4.5 mL/min. For each element, the calibration curves (Figure 1) were

Table 1 The instrumental settings for the ICP-MS

| | |
|---------------------------------|---------------------------------------|
| Spray chamber: | Quartz Cyclonic |
| Nebulizer: | Meinhard® Concentric Quartz Type A0.5 |
| Torch: | Standard Quartz |
| Sampler Cone: | Platinum |
| Skimmer Cone: | Platinum |
| Plasma gas flow: | 18 L min^{-1} |
| Auxiliary gas flow: | 1.20 L min^{-1} |
| Nebulizer gas flow: | 1.04 L min^{-1} |
| ICP RF power: | 1550 W |
| Pulse voltage: | 900 V |
| Analog stage voltage: | -1637 V |
| KED mode cell entrance voltage: | -4 V |
| KED mode cell exit voltage: | -38 V |
| KED mode axial field voltage: | 475 V |




Figure 1 Calibration curves for each element.

achieved using five standard concentrations (0, 0.1, 1, 10 and 100 µg/L) in 1% nitric acid solutions.

QC samples with concentrations of 0.5, 5 and 50 µg/L were analysed after calibration and at the end of samples determination. A 10 µg/L multielement standard was inserted in the analytical sequence every five samples for CCV. Three replicas were recorded for each determination. All analytes were determined with a dwell time of 50 ms per AMU and 1000 ms integration time. Element concentrations were expressed as mg/kg (ppm) in the original samples.

Instrument detection limit (IDL) with 95% confidence level was established for each analyte by the three-time multiplication of the standard deviation obtained from the analysis of ten runs of blank samples in the same day with the determinations. We have obtained the following IDLs: ⁶³Cu, 0.015 µg/kg; ⁶⁶Zn, 0.049 µg/kg; ¹¹¹Cd, 0.019 µg/kg; ²⁰⁸Pb, 0.074 µg/kg.

Method detection limits (MDL) were estimated to be: ⁶³Cu, 0.129 µg/kg; ⁶⁶Zn, 0.264 µg/kg; ¹¹¹Cd, 0.192 µg/kg; ²⁰⁸Pb, 0.143 µg/kg. For quality assurance of plant analysis for metals, recovery tests were performed by

analysing spiked real samples. Average recoveries (n = 5) were: ⁶³Cu, 100%; ⁶⁶Zn, 109%; ¹¹¹Cd, 88%; ²⁰⁸Pb, 91.

Results and discussion

Statistics

We analysed the concentrations of heavy metals using descriptive statistics and non-parametric analysis using Mintab 16 (Minitab Inc.). Data normality was tested using Shapiro-Wilk test. For all statistical analyses we used $\alpha = 0.05$. First, we tested the differences between the concentrations of heavy metals in the four analysed species (i.e. *Ceratophyllum demersum*, *Potamogeton pectinatus*, *Potamogeton perfoliatus*, *Potamogeton lucens*) using ANOVA non-parametric Kruskal-Wallis test. Further, we grouped the species on genus and tested the differences of heavy metals concentration between *Ceratophyllum* and *Potamogeton* genera using Mann-Whitney test. In order to test whether the heavy metals concentration from all the analysed samples shows sampling site differences on the Bazias – Coronini sector, we used the same ANOVA non-parametric Kruskal-Wallis test. When Kruskal-Wallis tests indicated significant differences between the analysed

groups, the analysis was further developed using Dwass-Steel-Critchlow-Fligner test for multiple comparisons. Finally, we tested the associations between Zn and Cd elements concentration using Spearman rank correlation [27].

Metals concentration in water and plant samples

The concentration of heavy metals in the water samples we analysed is much lower than the Romanian standards for freshwater quality [28]. The concentration values (given as average of the results from the four campaigns) are shown in Table 2. Lead level is the only one that may generate a possible threat to the water quality in Coronini sampling point.

As a general observation, most of the measured concentrations show a great dispersion (high values of the standard deviation - see Tables 2, 3 and 4). This can be a consequence of the fact that the samples are collected over a longer period of time (18 months) and during different vegetation seasons. Annual balance of precipitation, extreme river conditions (flooding in the spring, drought in the summer) and biological activity of different intensity can also influence the metal uptake rate. For the cadmium in water samples, the high dispersion of data can be determined by the metal low level of concentration (the average value is 2 ng/L). Because of the high values of standard deviation, the data follow a non-normal distribution. Thus, we have chosen to analyse the data with non-parametric statistics.

Ceratophyllum demersum L. is a free floating perennial macrophyte growing in slow flow and nutrients-rich water bodies. It represents an important food source for invertebrates, fish and herbivorous aquatic birds. The analysed *Potamogeton* species (*Potamogeton perfoliatus* L., *Potamogeton lucens* L., and *Potamogeton pectinatus* L.) are submerged, rooted, flowering aquatic plants that grow in alkaline waters. They form beds that represent food source and shelter for fish species, aquatic invertebrates, waterfowls etc. Both genera contribute to the establishment of the protected habitat on European level - 3150 - Natural eutrophic lakes with *Magnopotamion* and *Hydrocharition* - type vegetation [29].

Table 2 Metal concentration (average ± 1 standard deviation) in water samples

| Sampling point | Cadmium (µg/L) | Copper (µg/L) | Lead (µg/L) | Zinc (µg/L) |
|-------------------|----------------|---------------|-------------|-------------|
| Baziaş | 0.002 ± 0.004 | 1.89 ± 0.04 | SLD | 1.82 ± 3.49 |
| Divici | SLD | 3.17 ± 1.86 | SLD | 1.35 ± 1.89 |
| Coronini | 0.008 ± 0.006 | 1.46 ± 0.59 | 2.76 ± 3.22 | 0.78 ± 0.86 |
| Romanian standard | 1 | 20 | 5 | 100 |

Maximum admitted concentrations by the Romanian Standard for freshwater are included.

Table 3 Metals concentrations (average ± 1 standard deviation) in aquatic plant samples

| Plant species | Cadmium (µg/g) | Copper (µg/g) | Lead (µg/g) | Zinc (µg/g) |
|-----------------------|----------------|---------------|-------------|---------------|
| <i>C. demersum</i> | 3.52±2.07 | 22.71±21.08 | 20.06±8.68 | 104.23±102.76 |
| <i>P. pectinatus</i> | 0.64±0.91 | 6.24±2.99 | 6.63±6.47 | 16.38±8.04 |
| <i>P. perfoliatus</i> | 1.88±1.49 | 13.14±4.87 | 13.32±19.88 | 57.96±95.39 |
| <i>P. lucens</i> | 0.97±0.59 | 9.80±2.08 | 1.51±1.05 | 15.63±5.65 |

The species were chosen based on their previous use as bioindicators in aquatic ecosystems and due to the fact they are one of the most frequent aquatic plant species from the Danube River ecosystems in the Iron Gates Natural Park. Metals concentrations in the four aquatic plants considered in our survey are displayed in Table 3.

When grouping the samples on plant genus (i.e. *Ceratophyllum* and *Potamogeton*) and applying the Mann-Whitney test, differences appear for all the elements (U = 214, p=0.008 for cadmium, U = 216, p=0.007 for copper, U = 223, p=0.003 for lead, U = 241, p<0.001 for zinc). *Ceratophyllum* concentrates higher amounts of elements compared to *Potamogeton* species, being a better accumulator (Table 4). Mazej and Germ [3] mention that copper and zinc distribute evenly throughout *Potamogeton nodosus*, while lead concentrate in higher amounts in the plant root. Subsequently, the differences between the two genera for copper and zinc cannot be assigned to differences in elements uptake from the water body. Fritioff et al. [19] attribute lower metal concentration to species with higher biomass such as *Potamogeton*.

Differences also appear between the four species with the exception of *Ceratophyllum demersum* vs. *Potamogeton perfoliatus*, where they are not statistically significant (Dwass-Steel-Christlow-Fligner test = -0.73 p=0.96 for cadmium, (Dwass-Steel-Christlow-Fligner test = -0.51, p=0.98 for copper, Dwass-Steel-Christlow-Fligner test = -0.87, p=0.93 for lead and Dwass-Steel-Christlow-Fligner test = -1.82, p=0.57 for zinc). Our results, expressed as mean concentration ± 1 standard deviation for each species, indicate the highest concentration of all four elements in *Ceratophyllum demersum* (see Table 3). Among the *Potamogeton* species, the highest amounts of heavy metals are recorded in *Potamogeton perfoliatus* (1.88 µg/g for Cd; 13.14 µg/g for Cu; 13.32 µg/g for Pb; 57.96 µg/g for Zn). Our data are comparable with those obtained for cadmium and copper in the

Table 4 Metals concentrations (average ± 1 standard deviation) in *Ceratophyllum* and *Potamogeton* genera

| Plant genera | Cadmium (µg/g) | Copper (µg/g) | Lead (µg/g) | Zinc (µg/g) |
|----------------------|----------------|---------------|-------------|-------------|
| <i>Ceratophyllum</i> | 3.52±2.08 | 22.71±21.08 | 20.06±8.68 | 104.2±22.80 |
| <i>Potamogeton</i> | 1.42±1.38 | 10.75±5.18 | 12.26±16.19 | 41.60±6.10 |

same plant species by Pajevic et al. [25] on the same sector of the Danube River on the Serbian side.

We also grouped the plant samples according to the region they were collected from (with three delimited regions – Bazias, Divici, Coronini) and no differences appear between them (Kruskal-Wallis = 5.45, $p=0.07$ for Bazias; Kruskal-Wallis = 4.79, $p=0.09$ for Divici; Kruskal-Wallis = 3.22, $p=0.2$ for Coronini). The same conclusion was drawn for large-flow rivers by Fawzy et al. [30] for cadmium accumulation by plants (including *Ceratophyllum demersum*) from the Nile River ecosystems. In our case, the major pollution source is located near the Coronini sampling station and this is a 250 ha complex including settling ponds, sterile deposits and ores processing facilities from the former complex ores quarrying and processing plant. This had generated an increase of copper concentration in the sediments up to ten times between 1996 and 1999 (72 $\mu\text{g/g}$ and 679 $\mu\text{g/g}$, respectively) [26].

The long-term contamination of a water body leads to high amounts of metals in sediments which act like a deposit of heavy metals. Depending on the physical and chemical conditions and the water body regime (flooding, drought), they release metals into the water column, making them available for the aquatic vegetation. Metals removal from water by plants is a relatively rapid process [31]. Consequently, the metals are transferred to herbivorous fish species and aquatic invertebrates, thus entering the aquatic food chains and posing an ecotoxicological risk to species on higher trophic levels.

Starting from the hypothesis of a synergetic effect of zinc level on concentration of cadmium in the plants


[17], a Spearman test was applied to our set of data. The results confirm that cadmium concentration increases with zinc concentration (Spearman $\rho=0.40$, $p=0.02$). Since zinc is the heavy metal with the most important concentration in the region [26,32], it might enhance cadmium uptake by plants to significant levels, even if the ambient concentration is very low (0.002 $\mu\text{g/L}$).

Choosing the species to be used in such a survey is very important in a protected area, especially in a habitat of a Community importance. Since alien species introduction is to be avoided, the bioindicator species needs to be chosen from indigenous ones. Furthermore, they must be present on larger period of time during a year, in order to undertake exposure levels at different river regime (including extreme conditions such as drought and flooding). Nevertheless, they need to be present along the entire analysed sector, which was the case for *Trapa natans*, present, during the sampling period, only in Coronini. *Trapa natans* is also the species with the highest biomass and it is protected by the Habitats Directive [33].

At watershed scale, it is the first study on metals bioaccumulation by aquatic plants for the Romanian side of the Danube River. It allows comparison with the surveys performed for in Serbia [25,34].

Metal bioconcentration factors for *Ceratophyllum demersum*

The uptake rate of metals by plants from their surrounding environment is better described by the bioconcentration factor. For the floating species (*Ceratophyllum*



demersum L.), the bioconcentration factor was calculated as the ratio between the metal concentration in plant tissue and its concentration in water (as an average of the metal concentrations in the four sampling sites) [35].

The BCF values for *Ceratophyllum demersum* confirmed our hypothesis indicating the species as a hyper-accumulator for cadmium [14]. However, the values for the other elements are also high (see Figure 2). The sequence for the BCFs is Cd >> Zn > Pb > Cu.

The fact that the BCF value for cadmium is higher than 1000 ($1174.6 \text{ L} \times \text{g}^{-1}$) or 100-fold compared on a fresh weight [36] indicates *Ceratophyllum demersum* as a suitable species for phytoremediation purposes [37].

Conclusions

The ability to rapidly uptake the heavy metals from the environment, even at very low ambient levels, transforms the aquatic macrophytes into efficient indicators of the aquatic ecosystems quality. Differences appear between genera – with *Ceratophyllum* showing higher metals concentrations than *Potamogeton* – and species, with *Potamogeton perfoliatus* being the most effective accumulator of trace elements.

Furthermore, *Ceratophyllum demersum* has proven to be suitable for phytoremediation processes, since its capacity to hyperaccumulate cadmium from the water body reaches values of the bioconcentration factor higher than 1000. The negative aspect of their accumulation capacity is the fact the aquatic plants represent food sources for several categories of aquatic organisms, exposing them to contamination with toxic elements.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MLM designed the survey, collected the samples during the sampling campaigns and was in charge of data interpretation and article writing. CM was in charge of the analytical measurements (including quality control) and contributed to the article by writing the methods part. LR was responsible for the statistical processing and interpretation of data. AT was in charge of sample processing. All authors read and approved the final manuscript.

Acknowledgements

Dr. Marius Matache was supported by the strategic grant POSDRU/89/1.5/S/58852, Project "Postdoctoral programme for training scientific researchers" cofinanced by the European Social Fund within the Sectorial Operational Program Human Resources Development 2007 – 2013. This study was financed by the European Commission – Environment Directorate through the project LIFE10 NAT/RO/00740 "Improving the conservation status for the priority species and habitats in the Iron Gates wetlands".

Author details

¹University of Bucharest, Centre for Environmental Research and Impact Studies, Bucharest, Romania. ²"Emil Racoviță" Institute of Speleology of the Romanian Academy, Bucharest, Romania.

Received: 1 April 2013 Accepted: 25 September 2013

Published: 20 December 2013

References

1. Miretzky P, Saralegui A, Cirelli AF: Aquatic macrophytes potential for the simultaneous removal of heavy metals (Buenos Aires, Argentina). *Chemosphere* 2004, **57**:997–1005.
2. Dunbabin JS, Bowmer KH: Potential use of constructed wetlands for treatment of industrial wastewaters containing metals. *Sci Total Environ* 1992, **111**:151–168.
3. Mazej Z, Germ M: Trace element accumulation and distribution in four aquatic macrophytes. *Chemosphere* 2009, **74**:642–647.
4. Monterroso P, Abreu SN, Pereira E, Vale C, Duarte AC: Estimation of Cu, Cd and Hg transported by plankton from a contaminated area (Ria de Aveiro). *Acta Oecol* 200, **1**:351–357.
5. Albers PH, Camardese MB: Effects of acidification on metal accumulation by aquatic plants and invertebrates. 1. Constructed wetlands. *Environ Toxicol Chem* 1993, **12**:959–967.
6. Samecka-Cymerman A, Kempers AJ: Biomonitoring of Water Pollution with *Elodea Canadensis*. A Case Study of Three Small Polish Rivers with Different Levels of Pollution. *Water Air Soil Pollut* 2003, **145**:139–153.
7. Ostroumov SA, Shestakova TV: Decreasing the measurable concentrations of Cu, Zn, Cd, and Pb in the water of the experimental systems containing *Ceratophyllum demersum*: The phytoremediation potential. *Dokl Biol Sci* 2009, **428**:444–447.
8. Peng K, Luo C, Lou L, Li X, Shen Z: Bioaccumulation of heavy metals by the aquatic plants *Potamogeton pectinatus* L. and *Potamogeton malaiianus* Miq. and their potential use for contamination indicators and in wastewater treatment. *Sci Total Environment* 2008, **392**:22–29.
9. Yoon J, Cao X, Zhou Q, Ma LQ: Accumulation of Pb, Cu, and Zn in native plants growing on a contaminated Florida site. *Sci Total Environ* 2006, **368**:456–464.
10. Du Laing G, Rinklebe J, Vandecasteele B, Meers E, Tack FM: Trace metal behaviour in estuarine and riverine floodplain soils and sediments: a review. *Sci Total Environ* 2009, **407**:3972–3985.
11. Overesch M, Rinklebe J, Broll G, Neue HU: Metals and arsenic in soils and corresponding vegetation at Central Elbe river floodplains (Germany). *Environ Pollut* 2007, **145**:800–812.
12. Fritioff A, Greger M: Uptake and distribution of Zn, Cu, Cd, and Pb in an aquatic plant *Potamogeton natans*. *Chemosphere* 2006, **63**:220–227.
13. Vardanyan LG, Ingole BS: Studies on heavy metal accumulation in aquatic macrophytes from Sevan (Armenia) and Carambolim (India) lake systems. *Environ Int* 2006, **32**:208–218.
14. Rai UN, Sinha S, Tripathi RD, Chandra P: Wastewater treatability potential of some aquatic macrophytes: Removal of heavy metals. *Ecol Eng* 1995, **5**:5–12.
15. Estrella-Gómez N, Mendoza-Cózatl D, Moreno-Sánchez R, González-Mendoza D, Zapata-Pérez O, Martínez-Hernández A, Santamaría JM: The Pb-hyperaccumulator aquatic fern *Salvinia minima* Baker, responds to Pb²⁺ by increasing phytochelatin synthesis via changes in SmPCS expression and in phytochelatin synthase activity. *Aquat Toxicol* 2009, **91**:320–328.
16. Pell A, Marquez A, Lopez-Sanchez JF, Rubio R, Barbero M, Stegen S, Queirolo F, Diaz-Palma P: Occurrence of arsenic species in algae and freshwater plants of an extreme arid region in northern Chile, the Loa River Basin. *Chemosphere* 2013, **90**:556–564.
17. Bunluesin S, Pokethitayook P, Lanza GR, Tyson JF, Kruatrachue M, Xing B, Upatham S: Influences of Cadmium and Zinc Interaction and Humic Acid on Metal Accumulation in *Ceratophyllum Demersum*. *Water Air Soil Pollut* 2006, **180**:225–235.
18. Saygıdeger S, Dogan M: Lead and cadmium accumulation and toxicity in the presence of EDTA in *Lemna minor* L. and *Ceratophyllum demersum* L. *Bull Environ Contam Toxicol* 2004, **73**:182–189.
19. Fritioff A, Kautsky L, Greger M: Influence of temperature and salinity on heavy metal uptake by submersed plants. *Environ Pollut* 2005, **133**:265–274.
20. Weis JS, Weis P: Metal uptake, transport and release by wetland plants: implications for phytoremediation and restoration. *Environ Int* 2004, **30**:685–700.
21. Sommerwerk N, Baumgartner C, Bloesch J, Hein T, Ostojčić A, Paunović M, Schneider-Jacoby M, Siber R, Tockner K: Chapter 3 - The Danube River Basin. In *Rivers of Europe*. London: Academic Press; 2009:59–112.
22. Woitke P, Wellmütz J, Helm D, Kube P, Lepom P, Litheraty P: Analysis and assessment of heavy metal pollution in suspended solids and sediments of the river Danube. *Chemosphere* 2003, **51**:633–642.

23. Keller BEM, Lajtha K, Cristofor S: **Trace metal concentrations in the sediments and plants of the Danube Delta, Romania.** *Wetlands* 1998, **18**:42–50.
24. Tudor MI, Tudor M, David C, Teodorof L, Tudor D, Ibram O: **Heavy metals concentrations in aquatic environment and living organisms in the Danube Delta, Romania.** In *Chemicals as Intentional and Accidental Global Environmental Threats*. Edited by Simeonov L, Chirila E. The Netherlands: Springer; 2006:435–442. doi:10.1007/978-1-4020-5098-5_40.
25. Pajević S, Borišev M, Rončević S, Vukov D, Igić R: **Heavy metal accumulation of Danube river aquatic plants — indication of chemical contamination.** *Central European J Biol* 2008, **3**:285–294.
26. Matache M, Pătroescu C, Pătroescu-Klotz I: **Evoluția concentrației metalelor grele în sedimente acvifere dunărene pe tronsonul Baziaș-Portițele de Fier I (1996–1999).** *Rev Chim* 2002, **53**:623–626.
27. Zar JH: *Biostatistical Analysis. Fifth Edition edn.* Prentice Hall: Upper Saddle River, New Jersey; 2010.
28. MEWM: **Order of the Minister of Environment and Water Management no.161. for the approval of the norms regarding freshwater classification for establishing the ecological status of water bodies.** In *Book Order of the Minister of Environment and Water Management no.161. for the approval of the norms regarding freshwater classification for establishing the ecological status of water bodies*, Volume 511bis. City: Romanian Official Bulletin; 2006:2–92.
29. EC: **Council Directive 2006/105/EC of 20 November 2006 adapting Directives 73/239/EEC, 74/557/EEC and 2002/83/ EC in the field of environment, by reason of the accession of Bulgaria and Romania.** *Off J Eur Union* 2006, **363**:368–408.
30. Fawzy MA, Badr Nel S, El-Khatib A, Abo-El-Kassem A: **Heavy metal biomonitoring and phytoremediation potentialities of aquatic macrophytes in River Nile.** *Environ Monit Assess* 2012, **184**:1753–1771.
31. Forstner U, Wittmann GTW: *Heavy Metal Pollution in the Aquatic Environment.* 2nd edition. Berlin: Springer-Verlag; 1981.
32. Matache M, Rozyłowicz L, Ropota M, Patroescu C: **Heavy metals contamination of soils surrounding waste deposits in Romania.** *J Phys IV France* 2003, **107**:851–854.
33. Gafta D, Mountford JO (Eds.): *Manual de interpretare a habitatelor Natura 2000 din România [Romanian Manual for Interpretation of EU Habitats]*. Risoprint: Cluj-Napoca; 2008.
34. Stankovic Z, Pajevic S, Vuckovic M, Stojanovic S: **Concentration of trace metals in dominant aquatic plants of the Lake Provala (Vojvodina, Yugoslavia).** *Biol Plant* 2000, **43**:583–585.
35. Thiebaut G, Gross Y, Gierlinski P, Boiche A: **Accumulation of metals in *Elodea canadensis* and *Elodea nuttallii*: implications for plant-macroinvertebrate interactions.** *Sci Total Environ* 2010, **408**:5499–5505.
36. Zayed A, Gowthman S, Terry N: **Phytoaccumulation of trace elements by wetland plants: I Duckweed.** *J Environ Qual* 1998, **27**:715–721.
37. Phetsombat S, Kruatrachue M, Pokethitiyook P, Upatham S: **Toxicity and bioaccumulation of cadmium and lead in *Salvinia cucullata*.** *J Environ Biol* 2006, **27**:645–652.

doi:10.1186/2052-336X-11-39

Cite this article as: Matache et al.: Plants accumulating heavy metals in the Danube River wetlands. *Journal of Environmental Health Sciences & Engineering* 2013 **11**:39.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

